PubMed Services

Journals Database

MeSH Database Single Citation Matcher Baich Citation Matcher

Clinical Queries Special Queries LinkOut

Related Resources

Order Documents NLM Mobile

**NLM Catalog** 

TOXNET Consumer Health

NLM Gateway

Clinical Alerts

ClinicalTrials.cov

PubMed Central

My NCBI





### A service of the National Library of Medicine and the National Institutes of Health

My NCBI [7] [Sign In] [Register]

All Databases	PubMed	Nucleotide	Protein	Genome	Structure	OMIM	PMC	Journals	Books
Search PubMed	for polyethylene glycol and lactulose Go Clear								
	/ Limits	Preview/Index	→ History	Clipboard	> Details	*			
About Entrez	Note:	Performing y	our origina	l search, <i>poly</i>	ethylene gl	ycol and	lactulose	, in PubMo	ed will
Text Version		retrieve <u>58 ci</u>	tations.						
Entrez PubMed Overview	Display 3	bstract	¥	Show 20	Sort by	<b>▼</b> Sen	d to ▼		
Help   FAQ Tutorials	All: 1 p	Review: 0 🔭							
New/Noteworthy E-Utilities	□ 1: <u>Dig</u>	<u>Dis Sci.</u> 1984	Sep;29(9):8	809-16.				Related Art	ticles, Links

Small intestinal permeability to mannitol, lactulose, and polyethylene glycol 400 in celiac disease.

## <u>Ukabam SO</u>, <u>Cooper BT</u>.

Mannitol (molecular weight 182), lactulose (342), and polyethylene glycol 400 (range 242-550) absorption was studied in 25 controls, 22 untreated celiacs, and 13 treated celiacs. Untreated celiacs absorbed less mannitol and more lactulose than controls. Absorption of higher as well as lower molecular-weight polyethylene glycols was reduced in untreated celiac disease. Absorption returned towards normal on treatment. Polyethylene glycol and lactulose absorption was enhanced by administering them in a hypertonic solution. Polyethylene glycol 400 but not lactulose or mannitol was lipid soluble in vitro. It was concluded that the mucosa in untreated celiac disease was more "leaky" than normal. Polyethylene glycol 400 absorption data suggested that its absorption may largely be determined by its lipid solubility and was decreased in celiac disease because of the reduced surface area of the small intestine. Polyethylene glycol 400 cannot be recommended as a suitable marker for permeability studies of the small intestine.

PMID: 6432500 [PubMed - indexed for MEDLINE]

Display Abstract Show 20 Sort by Send to	

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Aug 14 2006 08:07:58





#### A service of the National Library of Medicine and the National Institutes of Health

My NCBI [Sign In] [Register]

All Databases	PubMed	Nucleotide	Protein	Genome	Structure	OMIM	FMC	Journals	Books
Search PubMed	•		for polyethylene glycol and lactulose						
	Limits	Preview/Index	→ History	Clipboard	> Details	×			

About Entrez

Text Version

E-Utilities


My NCBI

Entrez PubMed Overview Help | FAQ Tutorials New/Noteworthy

**PubMed Services** Journais Database MeSH Database Single Citation Matcher **Baich Citation Matcher** Clinical Queries Special Queries LinkOut

Related Resources Order Documents NLM Mobile NLM Catalog NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

Performing your original search, polyethylene glycol and lactulose, in PubMed will retrieve 58 citations.



1: Clin Pediatr (Phila). 2002 May;41(4):225-9.

Related Articles, Links



Comparison of polyethylene glycol 3350 and lactulose for treatment of chronic constipation in children.

## Gremse DA, Hixon J, Crutchfield A.

The Division of Pediatric Gastroenterology and Nutrition, University of South Alabama College of Medicine, Mobile 36604, USA.

Polyethylene glycol (PEG) 3350 and lactulose were compared in an unblinded, randomized, crossover design for treatment of constipation in 37 children aged 2 to 16 years. Subjects received lactulose (1.3 g/kg/d divided twice daily up to 20 g) or PEG 3350 (10 g/m<sup>2</sup>/day) for 2 weeks. PEG 3350 significantly decreased the total colonic transit time compared to lactulose (47.6+/-2.7 vs 55.3+/-2.4 hours, mean +/- SE, PEG 3350 vs lactulose, respectively, p = 0.038). The stool frequency, form, and the ease of passage were similar for each laxative. Polyethylene glycol 3350 is an effective laxative for the treatment of chronic constipation in children.

# **Publication Types:**

- Clinical Trial
- Randomized Controlled Trial

PMID: 12041718 [PubMed - indexed for MEDLINE]

					•••••••••	
<del>}</del>	*****************	900000000000	***********	***********	**********	
Diaming Abstract		32A 3	CAR!	h., ::::::::::::::::::::::::::::::::::::	Cand to	332233
1 )ISDIAV JADOUAUL	Stow	120	o <del>z</del> ogg OUI €	by 💮 🔀	Send to	333
210014	20000000 10110 11	4 9	*************		ė –	***************************************

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

Aug 14 2006 08:07:58